

The very-long-chain hydroxy fatty acyl-CoA dehydratase PASTICCINO2 is essential and limiting for plant development

Liên Bach*, Louise V. Michaelson[†], Richard Haslam[†], Yannick Bellec*, Lionel Gissot^{**}, Jessica Marion*, Marco Da Costa^{*§}, Jean-Pierre Boutin[¶], Martine Miquel[¶], Frédérique Tellier[¶], Frederic Domergue^{**}, Jonathan E. Markham^{††}, Frederic Beaudoin[†], Johnathan A. Napier[†], and Jean-Denis Faure^{**‡}

*Laboratoire Biologie Cellulaire, [†]Laboratoire Biologie des Semences, [‡]Plateforme de Cytologie et d'Imagerie Végétale, and [¶]Plateforme de Chimie du Végétal, Institut National de la Recherche Agronomique, 78000 Versailles Cedex, France; [§]Rothamsted Research, Harpenden, Herts AL5 2JQ, United Kingdom; ^{**}Laboratoire Biogenèse Membranaire, Centre National de la Recherche Scientifique-Université Bordeaux 2, BP 33076 Bordeaux Cedex, France; and ^{††}Donald Danforth Plant Science Center, Saint Louis, MO 63132

Edited by Roland Douce, Université de Grenoble, Grenoble, France, and approved July 14, 2008 (received for review May 27, 2008)

Very-long-chain fatty acids (VLCFAs) are synthesized as acyl-CoAs by the endoplasmic reticulum-localized elongase multiprotein complex. Two *Arabidopsis* genes are putative homologues of the recently identified yeast 3-hydroxy-acyl-CoA dehydratase (*PHS1*), the third enzyme of the elongase complex. We showed that *Arabidopsis* PASTICCINO2 (*PAS2*) was able to restore *phs1* cytokinesis defects and sphingolipid long chain base overaccumulation. Conversely, the expression of *PHS1* was able to complement the developmental defects and the accumulation of long chain bases of the *pas2-1* mutant. The *pas2-1* mutant was characterized by a general reduction of VLCFA pools in seed storage triacylglycerols, cuticular waxes, and complex sphingolipids. Most strikingly, the defective elongation cycle resulted in the accumulation of 3-hydroxy-acyl-CoA intermediates, indicating premature termination of fatty acid elongation and confirming the role of *PAS2* in this process. We demonstrated by *in vivo* bimolecular fluorescence complementation that *PAS2* was specifically associated in the endoplasmic reticulum with the enoyl-CoA reductase CER10, the fourth enzyme of the elongase complex. Finally, complete loss of *PAS2* function is embryo lethal, and the ectopic expression of *PHS1* led to enhanced levels of VLCFAs associated with severe developmental defects. Altogether these results demonstrate that the plant 3-hydroxy-acyl-CoA dehydratase PASTICCINO2 is an essential and limiting enzyme in VLCFA synthesis but also that *PAS2*-derived VLCFA homeostasis is required for specific developmental processes.

cuticular wax | elongase | sphingolipid | triacylglycerol | leaf development

Very-long-chain fatty acids (VLCFAs) are components of eukaryotic cells and are composed of 20 or more carbons (i.e., >C18). VLCFAs are involved in many different physiological functions in different organisms. They are abundant constituents of some tissues like the brain (myelin) or plant seeds (storage triacylglycerols). VLCFAs are components of the lipid barrier of the skin and the plant cuticular waxes (1). VLCFAs are also involved in the secretory pathway for protein trafficking and for the synthesis of GPI lipid anchor (2). Finally, VLCFAs are components of sphingolipids that are both membrane constituents and signaling molecules (3).

In yeast, VLCFA synthesis is catalyzed in the endoplasmic reticulum (ER) by a membrane-bound multienzyme protein complex referred as the elongase (4). The elongase complex catalyzes the cyclic addition of a C₂-moiety obtained from malonyl-CoA to an acyl-CoA. VLCFAs (C₂₀, C₂₂, C₂₄, or higher) are produced from shorter fatty acids (usually C₁₆ or C₁₈) made by the cytosolic fatty acid synthase complex. The two-carbon addition during the elongation cycle requires four independent but sequential enzymatic steps. The first step involves the condensation of the malonyl-CoA with an acyl-CoA precursor, resulting in a 3-ketoacyl-CoA intermediate,

which is reduced to form a 3-hydroxy-acyl-CoA. The third enzymatic step is the dehydration of the 3-hydroxy-acyl-CoA to an enoyl-CoA, which is finally reduced to yield an acyl_{*n* + 2}-CoA. The keto and enoyl reductases are encoded respectively by the YBR159w and TSC13/YDL015c genes; the condensing enzymes are coded by a small family of genes, *ELO1*, 2, and 3, of which *ELO2/FEN1/YCR034w* and *ELO3/SUR4/YLR372w* have been shown to be required for the synthesis of VLCFAs (5). Identification of the dehydratase remained elusive until the recent identification of YJL097w/*PHS1* as encoding this activity (although a role in sphingolipid biosynthesis had previously been inferred from the biochemical phenotype of *phs1* mutant that accumulated the long chain base phytosphingosine [PHS]) (6). The *phs1* mutant was also characterized as a cell cycle mutant defective in G₂/M phase (7). Ultimate confirmation of the biochemical function of Phs1p as the elongase dehydratase was provided by *in vitro* activity of recombinant protein and reconstitution of the elongase complex in proteoliposomes (8). Very recently, topology experiments demonstrated that Phs1p has six transmembrane domains with its N- and C-termini in the cytosol and that two conserved amino acids, Y149 and E152, were critical for its activity (9).

In plants, there is a large family of 3-ketoacyl-CoA synthases (KCS) condensing enzymes exemplified by the *Arabidopsis* gene *Fatty Acid Elongase 1 (FAEI)*, required in seeds for the synthesis of C₂₀+ fatty acids (e.g. erucic acid). The *Arabidopsis* genome encodes 21 FAE-like KCSs, and although these enzymes are structurally unrelated to the ELO class of condensing enzymes, it has been demonstrated that several *Arabidopsis* FAE-KCSs can rescue the otherwise lethal yeast *elo2Δ/elo3Δ* double mutant (10, 11). The *Arabidopsis* CER10 protein shows significant homology with yeast enoyl-CoA reductase Tsc13p, because it rescues the temperature-sensitive lethality of the *tscl3-1* yeast mutant (12) and is involved in VLCFA synthesis (13). The *Arabidopsis* genome also contains a gene that shares significant homology with the yeast 3-ketoreductase YBR159w, and although it has been demonstrated that this *Arabidopsis* activity

Author contributions: M.M., J.A.N., and J.-D.F. designed research; L.B., L.V.M., R.H., Y.B., L.G., J.M., M.D.C., J.-P.B., M.M., F.T., F.D., J.E.M., and F.B. performed research; J.-P.B. and J.E.M. contributed new reagents/analytic tools; F.D., J.A.N., and J.-D.F. analyzed data; and J.-D.F. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

[§]Present address: Laboratoire de Cytologie Expérimentale et Morphogenèse Végétale, Paris-VI, 94200 Ivry/Seine, France.

^{††}To whom correspondence should be addressed. E-mail: faure@versailles.inra.fr.

This article contains supporting information online at www.pnas.org/cgi/content/full/0805089105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA




Fig. 4. Acyl-CoA dehydratase is an essential and limiting activity. (A) Segregation of undeveloped seeds in *pas2-3/+* siliques (Bottom) compared with wild type (Top). (B) Ectopic expression of yeast *PHS1* (Right) impaired plant growth (Top) and modified leaf size and shapes (Bottom). (Scale bar, 3 cm at Top, 3 mm at Bottom.) (C) SEM of leaf epidermal cells of wild-type and *PHS1*-expressing plants (Top) and surface detail showing wax deposition in *PHS1* plants (Bottom). (Scale bar, 100 μ m at Top, 10 μ m at Bottom.) (D) Total fatty acid levels in the roots of *pas2-1-* and *PHS1*-expressing plants compared with wild type. (E) Total fatty acid levels in *pas2-1-* and *PHS1*-expressing plants compared with wild type. Dry weight and fatty acid values are the average of three samples \pm SD.

PHS1 expression on seed size and total fatty acid levels. Expression of *PHS1* led to slightly larger seeds, whereas *pas2* mutant showed smaller seeds compared with wild type (Fig. S11). Similar to the observation with seedlings, *PHS1*-expressing seeds showed an increase in VLCFAs, mostly 22:1 (Fig. 4E).

In conclusion, VLCFA dehydratase is not only an essential enzyme for plant growth and development, but it is also a limiting step for VLCFA synthesis because an increased dehydratase expression resulted in enhanced levels of VLCFAs in both vegetative and seed tissues.

Discussion

Collectively, our data demonstrated that the fatty acyl-CoA dehydratase is encoded by the *PASTICCINO2* gene, and that it is an essential activity for plants. Similarly, the ketoacyl-CoA reductase activity in the double *gl8agl8b* maize mutant was also found to be essential (24). However, the loss of function of the enoyl-CoA reductase *CER10*, which is involved in the ultimate step in the acyl-CoA elongation cycle, is not lethal (unlike the yeast orthologue TSC13), suggesting that there is at least another partially redundant *CER10* homologue in *Arabidopsis* (13). The weak *pas2-1* allele is still able to produce some very-long-chain acyl-CoAs, resulting in strong reduction but not complete absence of VLCFAs. However, the channeling of these rare VLCFAs into different lipid classes is not unselective in *pas2-1*, indicating a previously unobserved level of regulation for the homeostasis of different lipid types. The most straightforward explanation is that a leaky mutation such as *pas2-1* would result in significant perturbations to the lipid pools with the highest turnovers. Another hypothesis is that small changes in some lipid pools (like GIPCs) might lead to severe physiologic effects and that its homeostasis is maintained at the expense of less sensitive pools (like glucosylceramides). Alternatively, VLCFAs synthesis is compartmentalized differently and channeled independently for the different lipid pools. It was suggested, for instance, that VLCFA could be channeled into sphingolipids via an association

of ceramide synthases with the elongase complex. This hypothesis was raised to explain the accumulation of medium-chain ceramide observed in yeast elongase mutants as in *pas2* (25). It was previously thought that the limiting step in the elongase complex involved only the condensing enzymes (10). We demonstrated here that the dehydratase activity is also limiting for VLCFA synthesis. Interestingly, the overexpression of the condensing enzyme FAE1, as with *PHS1*, led to similar developmental alteration, such as asymmetric leaf shape. However, *PHS1* enhanced wax deposition whereas FAE1 suppressed it, suggesting that both enzymes, despite being part of the same complex, could modify VLCFA homeostasis in different ways (20).

Contrary to the *cer10* mutant, *pas2-1* showed very severe developmental defects. In particular, *pas2* was characterized by abnormal cell division that was enhanced in the presence of cytokinins, leading to callus-like structure (15, 16). It has to be noted that a similar phenotype was observed with weak mutations in the *PAS3/GURKE* gene, which codes the cytosolic acetylCoA carboxylase required for providing the malonyl-CoA to the elongase complex (26). The link between VLCFA and cell division was also reported in yeast (7, 27). In plants, the overexpression of *PAS2* delayed cell cycle progression, in particular during mitosis (19). *PAS2* was described as interacting directly with phosphorylated cell cycle regulator CDKA (19). The presence of a protein tyrosine phosphatase (PTP) motif that is conserved in eukaryotes led originally to the definition of the *PAS2/Phs1p* family as PTP-like proteins. However, recent structure/function analysis of *Phs1p* identified the catalytic residues involved in the dehydratase activity, and they do not belong to the PTP motif (9). We cannot exclude the possibility that the dehydratase function evolved recently from a PTP ancestor and that the PTP motif remained conserved across the *PAS2/Phs1p* family. Nonetheless, several indications suggest that the *PAS2/Phs1p* proteins might still be involved in phosphorylation-related processes. First, protein alignment of 31 members of *PAS2/Phs1p* showed amino acid conservation of the PTP motif (9). Then, the mutation of *PAS2/Phs1p* mammalian homolog PTPLA led to centronuclear myopathy in dogs, a disease related to mutations in the phosphoinositide phosphatase myotubularin MTM1 (28). *PHS1* had the strongest epistatic interaction in yeast with the LCB phosphatase *LCB3* (6). Moreover, the stability of the LCB kinase *LCB4p* is tightly regulated by the CDK PHO85p (29). The involvement of CDK-dependent phosphorylation in the regulation of LCB or VLCFA metabolic enzymes remains to be investigated in plants, but it would provide an attractive model reunifying the apparent divergent *PAS2* functions.

The nature of the *PAS2*-mediated VLCFA pathway that regulates cell division and cell differentiation is still unclear. Mutations downstream in the LCB and sphingolipid pathway will help in understanding the functional role of these different lipids in plant development. The fact that *PAS2* fulfills a nonredundant essential activity also opens up the possibility of using tissue-specific RNAi inactivation to probe and better define the multiple roles of VLCFAs in plant form and function.

Methods

Plant Material and Growth Conditions. The *pas2-1* mutants are ethyl methane sulfonate alleles in Col0 background that were maintained as heterozygous stocks. Plants were grown *in vitro* and in a greenhouse in soil as described previously (30). The pPAS2:*PAS2*-GFP construct corresponds to the *PAS2* genomic sequence with 1014 bp of promoter cloned into pMDC107. The *pas2-3* T-DNA insertion line N617051 from the SALK collection was genotyped by PCR with the *PAS2*-specific primers F20 (5'-AAAAAGCAGGCTC-GAGCTCGTCTAGTACACC-3') and R549 (5'-ACC CGGAAAATTCACAAATC-3') or T-DNA-specific primer Lba1 (5'-TGGTTCACGTAGTGGCCATCG-3'). Yeast strains and growth conditions were carried out as described previously (15).

Cytologic Analyses. Observations were carried out using an inverted TCS SP2-AOBS spectral confocal laser microscope (Leica Microsystems) using either

a PL APO 20 \times 0.70 NA or 63 \times 1.20 NA water-immersion objective. GFP and mCherry fluorescence were respectively recorded after an excitation at 488 and 594 nm (laser Ar/Kry) and a selective emission band of 495–550 nm and 600–643 nm. YFP fluorescence was recorded after an excitation at 514 nm (laser Ar/Kry) and a selective emission band of 520–564 nm. GUS (β -glucuronidase) staining and scanning electron microscopy were carried out as described previously (18). Colocalization and BiFC studies were performed as described previously (22).

Lipid Analyses. For mass quantification of lipid species, GIPCs, glucosylceramides, and ceramides from *A. thaliana* wild type and mutant seedlings were extracted, isolated, and quantified as detailed in (31). Analysis of free long

chain bases of sphingolipids was adapted from Lester and Dickson (32). Cuticular lipids were extracted and analyzed as described previously (33). Total seed and leaf fatty acid were analyzed as reported by Li *et al.* (34). AcylCoA profiling and MS/MS analysis were carried out as described previously (8, 35). Detailed description of lipid analysis can be found in the *SI Text*.

ACKNOWLEDGMENTS. We thank Patrick Moreau (Centre National de la Recherche Scientifique-Bordeaux 2), Olivier Grandjean, and Bruno Letarnec for their help and for discussion. J.M. and L.B. were funded, respectively, by 6th European Integrated Project AGRON-OMICS (grant no. LSHG-CT-2006–037704) and Cancéropôle Ile de France. Rothamsted Research receives grant-aided support from the Biotechnology and Biological Sciences Research Council (UK).

- Westerberg R, *et al.* (2004) Role for ELOVL3 and fatty acid chain length in development of hair and skin function. *J Biol Chem* 279:5621–5629.
- Gaigg B, Toulmay A, Schneiter R (2006) Very long-chain fatty acid-containing lipids rather than sphingolipids per se are required for raft association and stable surface transport of newly synthesized plasma membrane ATPase in yeast. *J Biol Chem* 281:34135–34145.
- Dickson RC, Sumanasekera C, Lester RL (2006) Functions and metabolism of sphingolipids in *Saccharomyces cerevisiae*. *Prog Lipid Res* 45:447–465.
- Tehlivets O, Scheuringer K, Kohlwein SD (2007) Fatty acid synthesis and elongation in yeast. *Biochim Biophys Acta* 1771:255–270.
- Kohlwein SD, *et al.* (2001) Tsc13p is required for fatty acid elongation and localizes to a novel structure at the nuclear-vacuolar interface in *Saccharomyces cerevisiae*. *Mol Cell Biol* 21:109–125.
- Schuldiner M, *et al.* (2005) Exploration of the function and organization of the yeast early secretory pathway through an epistatic miniarray profile. *Cell* 123:507–519.
- Yu L, Castillo LP, Mnaimneh S, Hughes TR, Brown GW (2006) A survey of essential gene function in the yeast cell division cycle. *Mol Biol Cell* 17:4736–4747.
- Denic V, Weissman JS (2007) A molecular caliper mechanism for determining very long-chain fatty acid length. *Cell* 130:663–677.
- Kihara A, Sakuraba H, Ikeda M, Denpoh A, Igarashi Y (2008) Membrane topology and essential amino acid residues of Phs1, a 3-hydroxyacyl-CoA dehydratase involved in very long-chain fatty acid elongation. *J Biol Chem* 283:11199–11209.
- Millar AA, Kunst L (1997) Very-long-chain fatty acid biosynthesis is controlled through the expression and specificity of the condensing enzyme. *Plant J* 12:121–131.
- Paul S, *et al.* (2006) Members of the *Arabidopsis* FAE1-like 3-ketoacyl-CoA synthase gene family substitute for the Elop proteins of *Saccharomyces cerevisiae*. *J Biol Chem* 281:9018–9029.
- Gable K, Garton S, Napier JA, Dunn TM (2004) Functional characterization of the *Arabidopsis thaliana* orthologue of Tsc13p, the enoyl reductase of the yeast microsomal fatty acid elongating system. *J Exp Bot* 55:543–545.
- Zheng H, Rowland O, Kunst L (2005) Disruptions of the *Arabidopsis* Enoyl-CoA reductase gene reveal an essential role for very-long-chain fatty acid synthesis in cell expansion during plant morphogenesis. *Plant Cell* 17:1467–1481.
- Dunn TM, Lynch DV, Michaelson LV, Napier JA (2004) A post-genomic approach to understanding sphingolipid metabolism in *Arabidopsis thaliana*. *Ann Bot (Lond)* 93:483–497.
- Bellec Y, *et al.* (2002) Pasticcino2 is a protein tyrosine phosphatase-like involved in cell proliferation and differentiation in *Arabidopsis*. *Plant J* 32:713–722.
- Faure JD, *et al.* (1998) The PASTICCINO genes of *Arabidopsis thaliana* are involved in the control of cell division and differentiation. *Development* 125:909–918.
- Haberer G, Erschadi S, Torres-Ruiz RA (2002) The *Arabidopsis* gene PEPINO/PASTICCINO2 is required for proliferation control of meristematic and non-meristematic cells and encodes a putative anti-phosphatase. *Dev Genes Evol* 212:542–550.
- Harrar Y, Bellec Y, Bellini C, Faure JD (2003) Hormonal control of cell proliferation requires PASTICCINO genes. *Plant Physiol* 132:1217–1227.
- Da Costa M, *et al.* (2006) *Arabidopsis* PASTICCINO2 is an antiphosphatase involved in regulation of cyclin-dependent kinase A. *Plant Cell* 18:1426–1437.
- Millar AA, Wrischer M, Kunst L (1998) Accumulation of very-long-chain fatty acids in membrane glycerolipids is associated with dramatic alterations in plant morphology. *Plant Cell* 10:1889–1902.
- Joubes J, *et al.* (2008) The VLCFA elongase gene family in *Arabidopsis thaliana*: Phylogenetic analysis, 3D modelling and expression profiling. *Plant Mol Biol* 67:547–566.
- Marion J, *et al.* (in press) Systematic analysis of protein subcellular localization and interaction using high-throughput transient transformation of *Arabidopsis* seedlings. *Plant J*.
- Tsegaye Y, *et al.* (2007) *Arabidopsis* mutants lacking long chain base phosphate lyase are fumonisin-sensitive and accumulate trihydroxy-18:1 long chain base phosphate. *J Biol Chem* 282:28195–28206.
- Dietrich CR, *et al.* (2005) Characterization of two GL8 paralogs reveals that the 3-ketoacyl reductase component of fatty acid elongase is essential for maize (*Zea mays* L.) development. *Plant J* 42:844–861.
- Han G, *et al.* (2002) The *Saccharomyces cerevisiae* YBR159w gene encodes the 3-ketoreductase of the microsomal fatty acid elongase. *J Biol Chem* 277:35440–35449.
- Baud S, *et al.* (2004) gurke and pasticcino3 mutants affected in embryo development are impaired in acetyl-CoA carboxylase. *EMBO Rep* 5:515–520.
- Al-Feel W, DeMar JC, Wakil SJ (2003) A *Saccharomyces cerevisiae* mutant strain defective in acetyl-CoA carboxylase arrests at the G2/M phase of the cell cycle. *Proc Natl Acad Sci USA* 100:3095–3100.
- Pele M, Turet L, Kessler JL, Blot S, Panthier JJ (2005) SINE exonic insertion in the PTPLA gene leads to multiple splicing defects and segregates with the autosomal recessive centronuclear myopathy in dogs. *Hum Mol Genet* 14:1417–1427.
- Iwaki S, Kihara A, Sano T, Igarashi Y (2005) Phosphorylation by Pho85 cyclin-dependent kinase acts as a signal for the down-regulation of the yeast sphingoid long-chain base kinase Lcb4 during the stationary phase. *J Biol Chem* 280:6520–6527.
- Smyczynski C, *et al.* (2006) The C terminus of the immunophilin PASTICCINO1 is required for plant development and for interaction with a NAC-like transcription factor. *J Biol Chem* 281:25475–25484.
- Markham JE, Jaworski JG (2007) Rapid measurement of sphingolipids from *Arabidopsis thaliana* by reversed-phase high-performance liquid chromatography coupled to electrospray ionization tandem mass spectrometry. *Rapid Commun Mass Spectrom* 21:1304–1314.
- Lester RL, Dickson RC (2001) High-performance liquid chromatography analysis of molecular species of sphingolipid-related long chain bases and long chain base phosphates in *Saccharomyces cerevisiae* after derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. *Anal Biochem* 298:283–292.
- Raffaele S, *et al.* (2008) A MYB transcription factor regulates very-long-chain fatty acid biosynthesis for activation of the hypersensitive cell death response in *Arabidopsis*. *Plant Cell* 20:752–767.
- Li Y, Beisson F, Pollard M, Ohlrogge J (2006) Oil content of *Arabidopsis* seeds: The influence of seed anatomy, light and plant-to-plant variation. *Phytochemistry* 67:904–915.
- Sayanova O, *et al.* (2006) A bifunctional $\Delta 12$, $\Delta 15$ -desaturase from *Acanthamoeba castellanii* directs the synthesis of highly unusual n-1 series unsaturated fatty acids. *J Biol Chem* 281:36533–36541.